

Patient care and clinical outcomes for patients with COVID-19 infection admitted to African high-care or intensive care units (ACCCOS): a multicentre, prospective, observational cohort study



The African COVID-19 Critical Care Outcomes Study (ACCCOS) Investigators*

Summary

Background There have been insufficient data for African patients with COVID-19 who are critically ill. The African COVID-19 Critical Care Outcomes Study (ACCCOS) aimed to determine which resources, comorbidities, and critical care interventions are associated with mortality in this patient population.

Methods The ACCCOS study was a multicentre, prospective, observational cohort study in adults (aged 18 years or older) with suspected or confirmed COVID-19 infection who were referred to intensive care or high-care units in 64 hospitals in ten African countries (ie, Egypt, Ethiopia, Ghana, Kenya, Libya, Malawi, Mozambique, Niger, Nigeria, and South Africa). The primary outcome was in-hospital mortality censored at 30 days. We studied the factors (ie, human and facility resources, patient comorbidities, and critical care interventions) that were associated with mortality in these adult patients. This study is registered on ClinicalTrials.gov, NCT04367207.

Findings From May to December, 2020, 6779 patients were referred to critical care. Of these, 3752 (55·3%) patients were admitted and 3140 (83·7%) patients from 64 hospitals in ten countries participated (mean age 55·6 years; 1890 [60·6%] of 3118 participants were male). The hospitals had a median of two intensivists (IQR 1–4) and pulse oximetry was available to all patients in 49 (86%) of 57 sites. In-hospital mortality within 30 days of admission was 48·2% (95% CI 46·4–50·0; 1483 of 3077 patients). Factors that were independently associated with mortality were increasing age per year (odds ratio 1·03; 1·02–1·04); HIV/AIDS (1·91; 1·31–2·79); diabetes (1·25; 1·01–1·56); chronic liver disease (3·48; 1·48–8·18); chronic kidney disease (1·89; 1·28–2·78); delay in admission due to a shortage of resources (2·14; 1·42–3·22); quick sequential organ failure assessment score at admission (for one factor [1·44; 1·01–2·04], for two factors [2·0; 1·33–2·99], and for three factors [3·66; 2·12–6·33]); respiratory support (high flow oxygenation [2·72; 1·46–5·08]; continuous positive airway pressure [3·93; 2·13–7·26]; invasive mechanical ventilation [15·27; 8·51–27·37]); cardiorespiratory arrest within 24 h of admission (4·43; 2·25–8·73); and vasopressor requirements (3·67; 2·77–4·86). Steroid therapy was associated with survival (0·55; 0·37–0·81). There was no difference in outcome associated with female sex (0·86; 0·69–1·06).

Interpretation Mortality in critically ill patients with COVID-19 is higher in African countries than reported from studies done in Asia, Europe, North America, and South America. Increased mortality was associated with insufficient critical care resources, as well as the comorbidities of HIV/AIDS, diabetes, chronic liver disease, and kidney disease, and severity of organ dysfunction at admission.

Funding The ACCCOS was partially supported by a grant from the Critical Care Society of Southern Africa.

Copyright © 2021 Elsevier Ltd. All rights reserved.

Introduction

SARS-CoV-2 has overwhelmed health-care systems by causing high rates of critical illness. The global case fatality rate of COVID-19 is approximately 3%,¹ with older people (eg, people older than 62 years) who have comorbidities known to be more susceptible than younger people.² Moreover, there is a concern of further mortality with subsequent waves across regions globally.

Our hypothesis was that critically ill patients with COVID-19 might have worse health outcomes in Africa than other continents because the ability to provide sufficient care is compromised by having a small

workforce,³ having a low number of intensive care facilities, and the scarcity of critical care resources.⁴ We also hypothesised that unplanned admissions would further adversely affect critical care outcomes in Africa⁵ as the ability of health-care systems to respond to meet the clinical workload is limited. Finally, patient outcomes following critical care for COVID-19 were not sufficiently documented in this under-resourced environment,⁶ despite a call for prevention and response measures in low-income and middle-income countries.⁷

As there were little data for the management of critically ill patients with COVID-19,⁸ we designed the African

Lancet 2021; 397: 1885–94

See [Comment](#) page 1862

*ACCCOS Investigators are listed in the appendix (pp 2–12)

Correspondence to:
Prof Bruce Biccard, Department of Anaesthesia and Perioperative Medicine, University of Cape Town, Cape Town 7925, South Africa; Department of Anaesthesia and Perioperative Medicine, Groote Schuur Hospital, Cape Town 7935, South Africa
bruce.biccard@uct.ac.za

See Online for appendix

Research in context

Evidence before this study

We searched MEDLINE, Embase, the Cochrane Library, Africa-Wide Information, and SciELO Citation Index between Jan 1 and Sept 23, 2020 (as well as an updated search from Sept 23 to Dec 6, 2020) using the search terms "(Betacoronavirus OR Betacoronaviruses)", "(Corona Virus OR Corona Viruses OR Coronavirus OR Coronaviruses)", "(COVID OR COVID19 OR COVID-19)", "(CoV OR CoV2 OR HCoV-19 OR nCoV OR 2019nCoV)", "(Severe Acute Respiratory Syndrome CoV OR severe acute respiratory syndrome coronavirus 2 OR SARS CoV 2 OR SARS-CoV-2 OR SARSCoV OR SARS-CoV OR SARS2)", "(Intensive care OR intensive care unit* OR ICU*)", "(ITU* or intensive treatment* OR intensive treatment unit*)", "critical care", "critical* ill*", "1 or 2 or 3 or 4 or 5", "6 or 7 or 8 or 9", "10 and 11", "Limit 12 to yr="2020". Studies that included patients with COVID-19 who were not in critical care, and critical care cohorts restricted to a specific patient subgroup were excluded. Studies published in all languages were considered in our search. There is little data to guide the management of critically ill patients with COVID-19 in under-resourced environments. Previously published systematic reviews confirmed that there were no published outcomes data from Africa, and little data for factors associated with mortality or survival in under-resourced environments.

Added value of this study

Mortality following critical care admission for patients with suspected COVID-19 infection and confirmed COVID-19 infection in this African cohort was 48.2% (95% CI 46.4–50.0). The meta-analysis reports a global mortality of 31.5% (27.5–35.5), with the African data reporting an excess mortality of 11 (in the best case scenario) to 23 (in the worst case

scenario) deaths per 100 patients compared with the global average. The excess mortality could be explained by the shortage of critical care resources. In our study, only one in two patients referred for critical care were admitted. Patients were admitted to units with limited access to dialysis, proning, extracorporeal membrane oxygenation (ECMO), arterial blood gases, and pulse oximetry. Furthermore, at the patient level, access to interventions (eg, dialysis, proning, and ECMO) were estimated to be between seven-times and 14-times lower than what is needed to manage critically ill patients with COVID-19. Adjusted analyses suggest that critical care mortality is associated with increasing age, the patient comorbidities of HIV/AIDS, diabetes, chronic liver disease, and kidney disease, the severity of organ dysfunction on presentation to critical care, and the initial need for increasing respiratory and cardiovascular support. The quick sequential organ failure assessment score at admission was associated with patient mortality and could be a simple and feasible risk stratification tool to use in under-resourced environments.

Implications of all the available evidence

In our study, mortality was strongly associated with organ dysfunction and the level of organ support needed at critical admission. The use of the quick SOFA score could provide guidance for appropriate triage decision making at the time of referral to critical care in an under-resourced setting when managing critically ill patients with COVID-19. Strategies are needed to mitigate risk in patients with COVID-19 in Africa with coexisting HIV/AIDS, diabetes, chronic liver disease, and kidney disease. It is likely that patient outcomes will continue to be severely compromised until the problems surrounding critical care resource scarcity are addressed.

COVID-19 Critical Care Outcomes Study (ACCCOS) to determine which resources, patient comorbidities, and critical care interventions were associated with mortality or survival in these patients. Wide dissemination of these findings could help to inform resource prioritisation necessary to manage patients who are critically ill with COVID-19 in Africa. This objective remains relevant as a recent meta-analysis⁶ reported no critical care outcomes data from Africa, or patient management data in resource-limited settings.

We aimed to determine which critical care resources, patient comorbidities, and hospital interventions were associated with in-hospital mortality in patients with suspected or confirmed COVID-19 who were referred to critical care in ten African countries.

Methods

Study design and participants

The ACCCOS study was a multicentre, prospective, observational cohort study in adults (aged 18 years or older) who were referred to intensive care or high-care

units with suspected or confirmed COVID-19 in ten African countries (ie, Egypt, Ethiopia, Ghana, Kenya, Libya, Malawi, Mozambique, Niger, Nigeria, and South Africa). The study was open to all African countries, and these ten countries fulfilled the ethics and regulatory requirements to participate. A high-care unit was defined as a patient area that provides a level of care between that given in an intensive care unit and a general ward but that does not usually provide invasive ventilation. Eligible patients included all patients admitted to a high-care or intensive care unit with suspected or confirmed COVID-19. Patient follow-up was until hospital discharge, censored at 30 days if the patient was in hospital. The study recruited from May 7 to Dec 18, 2020. The primary ethics approval was from the Human Research Ethics Committee of the University of Cape Town (Cape Town, South Africa).

All patients received standard of care for patients with suspected or confirmed COVID-19 and who required critical care admission. We planned to recruit as many sites as possible in Africa. Sites were requested to include

all eligible patients and to recruit for as long as possible with the understanding that they could stop recruiting at any point if they were overwhelmed by clinical commitments. Each site had to complete an eligible patients' screening log.

The primary ethics committee approved a so-called delayed consent process, as most patients would be unable to consent at the time of admission to critical care. The delayed consent process ensured consent by the patient (following stabilisation or recovery), a legal representative, or a proxy (in instances in which the patient was unable to provide consent). If there was no opportunity to acquire delayed consent before the study outcome was reached, then the ethics committee approved the inclusion of the patient's data in the study. The justification for this process was to minimise the risk of a non-consecutive patient enrolment, which would result in a biased sample. Some ethics committees waived consent. All sites fulfilled local ethics and regulatory requirements. The protocol and statistical analysis plan are posted at ClinicalTrials.gov, NCT04367207.

Procedures

Hospital specific data were collected by the hospital lead investigator before study registration and included: (1) the level of care, (2) the number of hospital and critical care beds, (3) reimbursement status, and (4) other factors affecting patient care (eg, nurse-to-patient ratio). The case record form and definitions document are in the appendix (pp 13–16).

To ensure a representative sample, we planned to include as many sites as possible with the requirement for inclusion of all consecutive patients using the delayed consent procedure.

Outcomes

The primary outcome of our study was in-hospital mortality within 30 days of admission. The secondary outcome was to determine the factors (ie, human and facility resources, patient comorbidities, and critical care interventions) that were associated with mortality in adult patients with suspected or confirmed COVID-19.

Statistical analysis

With 25–30 variables potentially associated with mortality in critically ill patients with COVID-19 (appendix pp 17–18) the minimum sample size required approximately 250–300 deaths to avoid violating the principle of approximately ten outcome events (ie, deaths) per variable in the regression.⁹ An interim analysis preprint¹⁰ was published in October, 2020, once that sample size was reached. The database was locked on Dec 18, 2020, with 1483 deaths in the cohort.

A statistical analysis plan was published on ClinicalTrials.gov before data evaluation for the interim analysis (appendix pp 17–18). Data were presented at an Africa wide level. Categorical variables were described as

proportions and were compared using χ^2 tests. Continuous variables were described as mean (SD) or median (IQR). Comparisons of continuous variables between groups were done using *t*-tests, one-way ANOVA, or equivalent non-parametric tests. The main model included patients with complete outcome data (ie, the main model excluded patients who were still in hospital and receiving therapy and who had not reached the outcome definition of death, discharge, or alive in hospital at 30 days). A three-level generalised linear mixed model (GLMM) was fitted using a logit link to identify independent risk factors for the primary outcome of mortality (with patients being at the first level, hospitals at the second level, and countries at the third level) to account for the expected correlation in outcomes within hospitals and countries. A fully conditional specification method was used to impute missing values for variables using an iterative Markov chain Monte Carlo method. We used a predictive mean matching method for scale variables. Five imputed datasets were constructed. All risk factors were considered for entry into the model provided there was no evidence of collinearity. The variables included subject variables, resource variables, and therapy variables. Subject variables included age, sex, body-mass index, coronary artery disease, congestive heart failure, hypertension, stroke or transient ischaemic attack, diabetes mellitus, cancer, whether the patient was a current smoker, chronic lung disease, active tuberculosis, chronic liver disease, HIV/AIDS, chronic or previous malaria, chronic kidney disease, cardiorespiratory arrest in the 24 h before referral to critical care, quick sequential organ failure assessment (SOFA) score,¹¹ and full SOFA¹² score on referral or admission. Resource variables included admission delayed due to the shortage of resources (eg, bed and staffing), nurse-to-patient ratio in critical care, ability to provide invasive ventilation, and physician availability on site 24 h per day, 7 days a week. Therapy variables included organ support at admission, respiratory support, proning, ventilatory support, intubation, inotropes or vasoconstrictors, dialysis, therapeutic anticoagulation, steroid therapy, repurposed or experimental COVID-19 drug therapy, and extracorporeal membrane oxygenation (ECMO). Collinearity was assessed using the variance inflation factor. Collinearity was associated with intubation, respiratory and ventilatory interventions, number of organs requiring support, and anticoagulation. Therefore, we created a single categorical variable for respiratory support (ie, none, oxygen, high flow nasal oxygen, continuous positive airway pressure, and invasive mechanical ventilation) and removed the dialysis and ECMO variables, which had collinearity with anticoagulation. The subsequent variance inflation factor showed collinearity between the subject variable “chronic malaria or malaria within 3 months” and the therapy variable “repurposed or experimental COVID-19

For the study protocol see https://clinicaltrials.gov/ProvidedDocs/07/NCT04367207/Prot_SAP_000.pdf

	Number of hospitals or median values
Hospital level of care	
Primary	4/57 (7%)
Secondary	13/57 (23%)
Tertiary	40/57 (70%)
Funding	
Government	50/57 (88%)
Private	3/57 (5%)
Dual funding	4/57 (7%)
Hospital type	
University affiliated	44/57 (77%)
Non-university affiliated	13/57 (23%)
Population served	2.0 million (0.2 million–5.0 million)
Hospital beds	500 (250–775)
Critical care beds providing invasive ventilation	12 (7–25)
Critical care beds unable to provide invasive ventilation	11 (8–29)
Specialist intensivists	2 (1–4)
Specialist doctors (not intensivists)	4 (2–8)
Non-specialist doctors	6 (2–11)
Nurse-to-patient ratio (day)	1:2 (1:2–1:1)
Nurse-to-patient ratio (night)	1:2 (1:3–1:1)
Doctor to patient ratio	1:4 (1:6–1:3)
Doctor on site in critical care after hours	49/57 (86%)
Surge capacity	
Number of extra intensive care ventilators	5 (3–10)
Number of operating room ventilators available for critical care upgrade	4 (1–8)
Haematology laboratory on hospital site	56/57 (98%)
Ability to do arterial blood gases on hospital site	45/55 (82%)
Critical care oxygen supply	
Vacuum insulated evaporator	27/57 (47%)
Cylinder oxygen	22/57 (39%)
Oxygen concentrator	8/57 (14%)
Pulse oximetry	
All patients	49/57 (86%)
Selected patients	7/57 (12%)
No patients	1/57 (2%)
Ability to provide prone ventilation	40/57 (70%)
Ability to provide renal replacement therapy	39/57 (68%)
Ability to provide extra-corporeal oxygenation	9/57 (16%)
Data are median (IQR) or n/N (%).	
Table 1: The characteristics of the hospitals included in the study	

drug therapy”. We removed “repurposed or experimental COVID-19 drug therapy” as this therapy variable was a heterogeneous variable compared with “chronic malaria or malaria within 3 months”. No further collinearity was identified. A three-level random-intercept mixed effects logistic regression was done on each of the five imputed




Figure: The ACCCOS recruitment

ACCCOS=African COVID-19 Critical Care Outcomes Study. *Three of 64 hospitals did not submit a screening log. The three hospitals without screening logs contributed 18 patients (ten patients from one hospital, six patients from another hospital, and two patients from a third hospital)

datasets using the `glmer` function in the `lme4` package¹³ in R.¹⁴ Estimates were combined from the five repeated complete data analyses using the `pool` function from the `mice` package.¹⁵ The `pool` function implements the rules for combining the separate estimates and SEs from each of the imputed datasets to provide an overall estimate with SEs, CIs, and p values.¹⁶ A p value of less than 0.05 was considered significant. To allow for comparison with the imputed datasets, the complete case analysis was also presented.

The results of the GLMM are reported as adjusted odds ratios with 95% CIs. Sensitivity analyses defined a priori were: (1) confirmed SARS-CoV-2-positive patients only, (2) confirmed SARS-CoV-2 patients excluding patients who had life support withdrawn or therapy limited (ie, the decision not to provide additional therapy, such as ventilation, adrenaline, and dialysis, to the patient's current therapy because of the expected poor prognosis), and (3) only patients who died or who were discharged alive (excluding in-hospital patients). A post-hoc sensitivity analysis that was requested by the reviewers excluded “cardiorespiratory arrest within 24 h of admission” as a potential variable. All analyses were done by AH and BMB.

A post-hoc decision was taken to update the critical care meta-analyses of COVID-19 mortality by region,^{6,8} done by EHT, KDMM, MELh, and JS. We have presented the case fatality rate for COVID-19 infections by region and did a meta-analysis of the mean age and SOFA score per region. The regional case fatality rate, critical care outcomes meta-analysis, and the meta-analysis of means for ages and SOFA scores provide context for rates of mortality in critical care in Africa.

Univariate analyses and the imputations were done using SPSS (version 26.0). The mixed effects logistic regressions were done using R.¹⁷ The meta-analyses were done using Stata 16 (StataCorp. 2019. *Stata Statistical Software*: Release 16. College Station, TX: StataCorp LLC). This study is registered on ClinicalTrials.gov, NCT04367207.

	All patients (n=3140)	Patients who died (n=1483)	Patients who survived (n=1594)	p value
Age, years				
Mean (SD)	56 (16.11)	58.74 (15.74)	52.68 (15.88)	<0.0001
Age quartiles, years				
18–45	832/3072 (27.1%)	302/1480 (20.4%)	530/1592 (33.3%)	<0.0001
46–56	713/3072 (23.2%)	317/1480 (21.4%)	396/1592 (24.9%)	..
57–67	769/3072 (25.0%)	409/1480 (27.6%)	360/1592 (22.6%)	..
>67	758/3072 (24.7%)	452/1480 (19.2%)	306/1592 (19.2%)	..
Sex				
Male	1890/3118 (60.6%)	902/1481 (60.9%)	966/1593 (60.6%)	0.883
Female	1228/3118 (39.4%)	579/1481 (39.1%)	627/1593 (39.4%)	..
Body-mass index categories				
<25	397/1596 (24.9%)	169/715 (23.6%)	221/851 (26.0%)	0.030
25–29.9	548/1596 (34.3%)	239/715 (33.4%)	300/851 (35.3%)	..
30–34.9	353/1596 (22.1%)	151/715 (21.1%)	198/851 (23.3%)	..
35–39.9	154/1596 (9.6%)	82/715 (11.5%)	65/851 (7.6%)	..
≥40	144/1596 (9.0%)	74/715 (10.3%)	67/851 (7.9%)	..
Missing	1544	768/1511 (50.8%)	743/1511 (49.2%)	..
Comorbidities				
Coronary artery disease	237/3093 (7.7%)	128/1469 (8.7%)	109/1586 (6.9%)	0.058
Congestive heart failure	209/3087 (6.8%)	100/1465 (6.8%)	106/1584 (6.7%)	0.885
Hypertension	1572/3104 (50.6%)	794/1476 (53.8%)	759/1588 (47.8%)	0.001
Stroke or transient ischaemic attack	139/3088 (4.5%)	77/1465 (5.3%)	62/1584 (3.9%)	0.082
Diabetes	1175/3090 (38.0%)	623/1471 (42.4%)	542/1582 (34.3%)	<0.0001
Cancer	96/3088 (3.1%)	49/1466 (3.3%)	45/1586 (2.8%)	0.463
Current smoker	140/3078 (4.5%)	56/1458 (3.8%)	80/1581 (5.1%)	0.114
Chronic lung disease	216/3088 (7.0%)	105/1465 (7.2%)	108/1586 (6.8%)	0.722
Active tuberculosis	51/3084 (1.7%)	30/1462 (2.1%)	20/1583 (1.3%)	0.116
Chronic liver disease	46/3086 (1.5%)	29/1465 (2.0%)	17/1585 (1.1%)	0.052
HIV/AIDS	237/3084 (7.7%)	149/1460 (10.2%)	86/1587 (5.4%)	<0.0001
Antiretroviral therapy	180/273 (65.9%)	115/175 (65.7%)	64/96 (66.7%)	0.894
Chronic malaria or malaria within 3 months	190/3089 (6.2%)	60/1465 (4.1%)	130/1586 (8.2%)	<0.0001
Chronic kidney disease	241/3085 (7.7%)	149/1465 (10.2%)	90/1585 (5.7%)	<0.0001
Condition at admission				
Cardiorespiratory arrest in 24 h before critical care referral	100/3086 (3.2%)	77/1467 (5.2%)	22/1583 (1.4%)	0.001
Organ support required at admission				
Respiratory support	2743/3102 (11.6%)	1375/1479 (93.0%)	1332/1587 (83.9%)	<0.0001
Cardiovascular support	609/296 (19.4%)	451/1396 (32.3%)	155/1537 (10.1%)	<0.0001
Renal support	282/3072 (9.2%)	185/1462 (12.7%)	95/1581 (6.0%)	<0.0001
Other support	510/3024 (16.9%)	217/1432 (15.2%)	293/1568 (18.7%)	0.011
Number of organ systems requiring support				
No organ system	271/3066 (8.8%)	92/1479 (6.2%)	179/1587 (11.3%)	<0.0001
One organ system	1816/3066 (59.2%)	797/1479 (53.9%)	1019/1587 (64.2%)	..
Two organ systems	705/3066 (23.0%)	389/1479 (26.3%)	316/1587 (19.9%)	..
Three organ systems	219/3066 (7.1%)	151/1479 (10.2%)	68/1587 (4.3%)	..
Four organ systems	55/3066 (1.8%)	50/1479 (3.4%)	5/1587 (0.3%)	..
Quick SOFA score on presentation				
SBP ≤100 mm Hg	471/3088 (15.3%)	326/1472 (22.1%)	141/1576 (8.9%)	<0.0001
Respiratory rate ≥22 breaths per min	2502/3085 (81.1%)	1237/1468 (84.3%)	1235/1578 (78.3%)	<0.0001
Glasgow Coma Scale score ≤14	850/3072 (27.7%)	588/1459 (40.3%)	253/1574 (16.1%)	<0.0001

(Table 2 continues on next page)

	All patients (n=3140)	Patients who died (n=1483)	Patients who survived (n=1594)	p value
(Continued from previous page)				
Quick SOFA score				
No risk factors	395/3069 (12.9%)	116/1458 (8.0%)	275/1573 (16.1%)	<0.0001
One risk factor	1786/3069 (58.2%)	744/1458 (51.0%)	1015/1573 (64.5%)	..
Two risk factors	641/3069 (20.9%)	395/1458 (27.1%)	240/1573 (15.3%)	..
Three risk factors	247/3069 (8.0%)	203/1458 (13.9%)	43/1573 (2.7%)	..
Full SOFA score on admission	4 (3–6)	5 (4–8)	4 (2–5)	<0.0001
Full SOFA score missing data	2136
Intensive care unit resources				
Admission delayed because of shortage of resources (eg, bed and staffing)	248/2947 (8.4%)	146/1382 (10.6%)	100/1530 (6.5%)	0.0001
Nurse-to-patient ratio	0.5 (0.25–1.00)	0.5 (0.33–1.00)	0.5 (0.25–0.67)	<0.0001
Ability to provide invasive ventilation for patient if required	2780/3091 (87.7%)	1361/1470 (92.6%)	1319/1585 (83.2%)	<0.0001
Physician available on site 24 h / 7 days a week	2760/3084 (89.5%)	1353/1463 (92.5%)	1376/1583 (86.9%)	<0.0001
Respiratory support (ie, highest level of support)				
None	421/2995 (14.1%)	237/1416 (16.7%)	182/1546 (11.8%)	<0.0001
Oxygen mask	1352/2995 (43.1%)	489/1416 (34.5%)	860/1546 (55.6%)	..
High-flow nasal oxygenation	589/2995 (19.7%)	293/1416 (20.7%)	274/1546 (17.7%)	..
CPAP	633/2995 (21.1%)	397/1416 (28.0%)	230/1546 (14.9%)	..
Prone ventilation				
None	1516/2826 (53.6%)	664/1305 (50.9%)	846/1503 (56.3%)	<0.0001
Not ventilated	989/2826 (35.0%)	396/1305 (30.3%)	581/1503 (38.7%)	..
Invasive mechanical ventilation	321/2826 (11.4%)	245/1305 (18.8%)	76/1503 (5.1%)	..
Ventilatory support				
None	1358/2902 (46.8%)	298/1384 (21.5%)	1053/1494 (70.5%)	<0.0001
Non-invasive ventilation	380/2902 (13.1%)	168/1384 (12.1%)	206/1494 (13.8%)	..
Invasive mechanical ventilation	1164/2902 (40.1%)	918/1384 (66.3%)	235/1494 (15.7%)	..
Intubation				
No	1896/3062 (61.9%)	532/1458 (36.5%)	1346/1576 (85.4%)	<0.0001
Yes (elective)	385/3062 (12.6%)	293/1458 (20.1%)	90/1576 (5.7%)	..
Yes (emergency)	781/3062 (25.5%)	633/1458 (43.4%)	140/1576 (8.9%)	..
Inotropes or vasoconstrictors	931/3086 (30.2%)	748/1472 (50.8%)	179/1587 (11.3%)	<0.0001
Dialysis	330/3073 (10.7%)	177/1467 (12.1%)	150/1579 (9.5%)	0.022
Therapeutic anticoagulation	2430/3084 (78.8%)	1218/1478 (82.4%)	1188/1579 (75.2%)	<0.0001
Steroid therapy	2519/3011 (83.7%)	1211/1430 (84.7%)	1281/1552 (82.5%)	0.125
Repurposed or experimental COVID-19 drug therapy	367/2937 (12.5%)	122/1367 (8.9%)	243/1549 (15.7%)	<0.0001
Extracorporeal membrane oxygenation (if available)	7/862 (0.8%)	5/359 (1.4%)	2/501 (0.4%)	0.110
Data are mean (SD), n (%), n (proportion), or median (IQR). Odds ratios were constructed for in-hospital mortality with univariate binary logistic regression analysis. CPAP=continuous positive airway pressure. SBP=systolic blood pressure. SOFA=sequential organ failure assessment.				
Table 2: Description of African COVID-19 critical care cohort				

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

Of the 40 African countries that were invited to participate via the African Perioperative Research Group (with 26 country leaders accepting the invitation), ten

countries (which included 64 hospitals) participated; Egypt (ten hospitals), Ethiopia (seven hospitals), Ghana (two hospitals), Kenya (three hospitals), Libya (14 hospitals), Malawi (three hospitals), Mozambique (two hospitals), Niger (two hospitals), Nigeria (eight hospitals), and South Africa (13 hospitals). 57 (89%) hospitals provided hospital level data (table 1). Most of the hospitals were university affiliated, government funded, and tertiary level hospitals. There was a median of

two intensivists (IQR 1–4), with a median nurse-to-patient ratio of 1:2 (1:2–1:1) during the day. 27 (47%) of 57 hospitals had oxygen available from vacuum insulated evaporators, with a median surge ventilator capacity of five intensive care ventilators (3–10) and four anaesthesia ventilators (1–8). Only 49 (86%) sites could provide pulse oximetry to all patients in critical care, and 39 (68%) could provide renal replacement therapy. 3752 (55·3%) of 6779 patients who were referred to critical care were admitted (figure). 3140 (83·7%) of 3752 patients who were eligible were included in our study, with a median of 25 patients (11–59) per hospital. The age range was 18 to 100 years, with men comprising 1890 (60·6%) of 3118 patients.

From May 7 to Dec 18, 2020, 6779 patients were referred to critical care (figure). 2995 (95·4%) of 3140 patients in the cohort were confirmed positive for SARS-CoV-2 infection. Patients were referred from the emergency department (937 [30·2%] of 3103 patients), from another department within the same hospital (926 [29·8%] patients), or another hospital (1240 [40·0%] patients). Mortality differed between the sites of referral ($p=0·002$), with in-hospital mortality of 401 (43·5%) of 921 patients from emergency departments, 443 (48·5%) of 914 patients from in-hospital referrals, and 629 (51·2%) of 1228 patients from other hospitals. Admission to critical care was delayed in 248 (8·4%) of 2947 patients because of a shortage of resources at the time of admission. A quick SOFA score of three had a mortality of 203 (82%) of 247 patients. Patients who required mechanical ventilation had a mortality rate of 78·9% (918 of 1164 patients).

The patients enrolled in our study had a mean age of 56 years with few comorbidities, and 39·4% were women (table 2). The most common comorbidities were hypertension, diabetes, HIV/AIDS, coronary artery disease, and chronic kidney disease. The quick SOFA score was completed in 3069 (97·7%) of 3140 patients at admission.¹¹ Most (2136 [68·0%] of 3140 patients) did not have a full SOFA score done at admission.

The length of critical care stay was 7 days (IQR 4–12). The decision to limit therapy was made in 284 (9·2%) of 3086 patients, and therapy was withdrawn in 81 (2·6%) patients. 72 (88·9%) of the 81 patients in whom therapy was withdrawn had already had therapy limited.

In-hospital mortality in the 30 days after admission to a high-care or critical care unit occurred in 1483 (48·2%; 95% CI 46·4–50·0) of 3077 patients, with 261 (16·4%) of 1594 patients alive and in hospital at 30 days (35 of these 244 [data from 17 patients is missing with respect to whether they were in an intensive care unit or not] patients were still in intensive care), and 1333 (83·6%) patients had been discharged. The primary outcome was unknown for 63 patients.

For secondary outcome measures, the missingness of data for the GLMM is given in the appendix (p 19). The GLMM for the risk factors associated with mortality

(ie, resources, comorbidities, and interventions) is shown in table 3. Risk factors that were independently associated with mortality were increasing age and the comorbidities of HIV/AIDS, diabetes, chronic liver disease or kidney disease, an increasing quick SOFA score, high flow nasal oxygenation, continuous positive airway pressure, invasive mechanical ventilation, and cardiorespiratory arrest within 24 h before admission to critical care, and the need for inotropes. Steroid therapy was associated with increased survival. The sex of the patient was not

	Odds ratio (97·5% CI)	p value
Patient characteristics		
Intercept	0·01 (0·0–0·04)	<0·0001
Age, years	1·03 (1·02–1·04)	<0·0001
Female	0·86 (0·69–1·06)	0·15
Body-mass index		
<25	1 (ref)	..
25·0–29·9	1·12 (0·78–1·61)	0·52
30·0–34·9	1·06 (0·73–1·53)	0·77
35·0–39·9	1·16 (0·74–1·81)	0·51
≥40	1·11 (0·66–1·86)	0·70
Comorbidities		
Coronary artery disease	0·80 (0·54–1·19)	0·27
Congestive heart failure	0·93 (0·63–1·38)	0·71
Hypertension	0·95 (0·76–1·19)	0·65
Stroke or transient ischaemic attack	1·04 (0·63–1·69)	0·89
Diabetes	1·25 (1·01–1·56)	0·04
Cancer	1·62 (0·90–2·91)	0·11
Current smoker	0·59 (0·35–1·01)	0·06
Chronic lung disease	0·96 (0·65–1·42)	0·85
Active tuberculosis	1·86 (0·85–4·06)	0·12
Chronic liver disease	3·48 (1·48–8·18)	0·004
HIV/AIDS	1·91 (1·31–2·79)	0·0008
Chronic malaria or malaria within 3 months	1·06 (0·52–2·15)	0·88
Chronic kidney disease	1·89 (1·28–2·78)	0·001
Cardiorespiratory arrest in 24 h before critical care referral	4·43 (2·25–8·73)	<0·0001
Quick SOFA score		
No risk factors	1 (ref)	..
One risk factor	1·44 (1·01–2·04)	0·04
Two risk factors	2·00 (1·33–2·99)	0·0008
Three risk factors	3·66 (2·12–6·33)	<0·0001
Intensive care unit resources		
Admission delayed because of shortage of resources (eg, bed and staffing)	2·14 (1·42–3·22)	0·0003
Physician available on site 24/7 for patient	0·84 (0·50–1·42)	0·51
Nurse-to-patient ratio (per unit increase)	1·31 (0·98–1·74)	0·07
Respiratory support		
None	1 (ref)	..
Oxygen mask	1·28 (0·74–2·20)	0·37
High flow nasal oxygenation	2·72 (1·46–5·08)	0·002
CPAP	3·93 (2·13–7·26)	<0·0001
Invasive mechanical ventilation	15·27 (8·51–27·37)	<0·0001

(Table 3 continues on next page)

	Odds ratio (97.5% CI)	p value
(Continued from previous page)		
Organ systems requiring support at admission		
No organ system	1 (ref)	..
One organ system	0.93 (0.56–1.52)	0.76
Two organ systems	0.93 (0.54–1.58)	0.78
Three organ systems	1.49 (0.76–2.93)	0.25
Four organ systems	2.99 (0.88–10.18)	0.08
Prone ventilation		
None	1 (ref)	..
On spontaneous ventilation	0.82 (0.61–1.09)	0.17
On invasive mechanical ventilation	1.42 (0.88–2.30)	0.15
Inotropes or vasoconstrictors	3.67 (2.77–4.86)	<0.0001
Therapeutic anticoagulation	1.18 (0.85–1.63)	0.33
Steroid therapy	0.55 (0.37–0.81)	0.003
CPAP=continuous positive airway pressure. SOFA=sequential organ failure assessment. Controls are patients alive in hospital and alive and discharged at 30 days (n=3140).		
Table 3: Generalised linear mixed model (pooled results of the imputed datasets) for patients referred to critical care with suspected or confirmed COVID-19 infection with full dataset for mortality		

associated with mortality. The sensitivity analyses supported these findings (appendix pp 20–25). No human resource factors (ie, physician available on site 24 h/7 days a week for patients and nurse-to-patient ratio) were associated with mortality.

A post-hoc analysis exploring the frequency of interventions received by patients who required invasive mechanical ventilation showed that 148 (12.9%) of 1146 of these patients also received dialysis.

The updated meta-analyses are shown in the appendix (pp 26–40). The overall reported global mortality was 31.5% (95% CI 27.5–35.5) and the mortality in Africa was higher than the mortality in the other regions. The excess mortality in Africa was 11 excess deaths per 100 admissions in Africa (in the best case scenario) and 23 excess deaths per 100 admissions in Africa (in the worst case scenario), compared with the global rate. The age and SOFA scores of this cohort were significantly lower than the global cohort, and the point estimates of this cohort were lower than the other regional cohorts (appendix pp 30–31). The case fatality rate in Africa was approximately 0.1% higher than in Europe and the global average, and similar to North America's case fatality rate. In Africa, up until Dec 11, 2020, there were 57 (40.1% of the global average) cases of COVID-19 per 100 000 population, which was lower than the global average of 140 cases per 100 000 population (appendix p 41).¹⁸

Discussion

The principal finding of our study was that in-hospital mortality following critical care admission for COVID-19 infection in Africa occurred in 48.2% (95% CI 46.4–50.0) of 3077 patients in the 30 days after high-care or intensive care unit admission, with an excess mortality of 11–23 deaths per 100 patients compared with the global

average. Mortality was associated with increasing age, HIV/AIDS, diabetes, chronic liver disease, kidney disease, a high severity of organ dysfunction on presentation, and increasing respiratory and cardiovascular support. A shortage of critical care resources could have contributed to increased mortality. The SOFA score could represent a simple, quick tool for risk stratification of patients with COVID-19 at the point of critical care admission.

Compared with other studies that have found men at higher risk of mortality with COVID-19, the finding of no sex differences in mortality in our study was unexpected.¹⁹ It is possible that women have an increased mortality risk generally because of the barriers to accessing care and limitations or biases in care when critically ill,²⁰ which could have moderated the differences between men and women here. Previously, the clinical course of patients with HIV and COVID-19 infection was unknown.²¹ Our data suggests that HIV/AIDS is an important risk factor for COVID-19 mortality. Our study also supports the use of steroid therapy to decrease mortality from COVID-19 in this patient population.²²

It is possible that overall mortality following SARS-CoV-2 infection is lower in Africa than in the other parts of the world (appendix p 41). However, the critical care in-hospital mortality was higher for African countries than for other, non-African countries. There are several possible reasons to explain this finding. First, the scarcity of critical care resources in African countries might contribute to the high critical care mortality. Second, there are inadequate critical care beds,⁴ with only one in two patients referred to critical care being admitted. Yet, the full SOFA scores suggest that the patients who were admitted could be relatively healthier than patients admitted in countries with more critical care resources (appendix p 31). Third, when considering the proportion of sites that could provide dialysis, proning, ECMO, arterial blood gases, and pulse oximetry, data from our study suggests that the African countries included in our study had very under-resourced critical care facilities. These data suggest that these countries had a low-volume critical care capacity, which could have adversely affected outcomes.²³ Lastly, merely counting the available critical care resources necessary for intervention does not accurately reflect the proportion of patients who actually receive the interventions. We estimate that patient access to interventions was between seven-times lower (for dialysis and proning) and 14-times lower (for ECMO) than what is required. Dialysis was available in 39 (68%) of 57 sites and was offered to only 330 (10%) of 3073 patients. Yet, acute kidney injury could occur in over 90% of patients with COVID-19 admitted to intensive care units, with one in four patients who have been ventilated requiring renal replacement therapy.²⁴

Proning is included in acute respiratory distress syndrome management strategies and has been provided in over 90% of patients.²⁵ As 75% of patients with COVID-19 referred to critical care develop acute

respiratory distress syndrome,²⁶ at least six times more patients should have received proning in our cohort. Similarly, ECMO was only available in nine (15·8%) of 57 sites (table 1), but ECMO was offered to less than 1% of patients; yet large registry data supports its use in patients with COVID-19 with refractory respiratory failure.²⁷ Lack of access to these interventions could partly explain the high mortality in Africa, and why one in eight patients had therapy withdrawn or limited.

The human resources available to these critical care units were somewhat good with respect to availability of a physician 24 h per day (7 days a week), and nurse-to-patient ratio. However, the inability to admit approximately half of the referred patients to the critical care unit could reflect intensive care units working at full capacity with the available resources, and therefore the effect of limited critical care human resources might have resulted in adverse outcomes outside the critical care unit, which we could not assess.

There are several limitations to our study. First, our study presents data from predominantly tertiary hospitals, yet pulse oximetry was not universally available. It is likely that lower level hospitals with less resourced critical care units might have had worse outcomes than those reported in this cohort. Furthermore, referral to higher level centres might have further increased mortality before patients were able to reach an appropriate critical care unit. We did not distinguish between cardiorespiratory arrests that occurred in hospital and those out of hospital, and it is likely that patients who had cardiorespiratory arrests outside of hospital might have had a higher mortality, and might be poorly represented in this cohort. It is therefore possible that the mortality for patients with COVID-19 who are critically ill might be higher than the mortality we report. The outcomes of the patients who were referred to critical care but not admitted are also unknown. It is unlikely that the findings of this study are generalisable to those patients, as their disease severity and resources available for therapy would differ from patients who are admitted to critical care.

Furthermore, we cannot report on the association between severity of comorbidities (eg, CD4 cell count in HIV/AIDS and mortality). Compared with other COVID-19 critical care cohorts (appendix p 30), our cohort was on average younger and it is therefore likely that the severity of comorbidities was less in this African cohort. Increasing age is associated with adverse outcomes in COVID-19 infections,²⁸ and it is therefore likely that our estimate of excess mortality is an underestimate when matched for age and severity of comorbidities.

The inability to recruit all eligible patients reflects the difficulties of doing research while providing a critical care service in an under-resourced environment during a pandemic. We have little data to help us to understand how one in two patients died without receiving oxygen, and how one in three patients died without receiving inotropes. It is unclear whether these events occurred

because of unavailability of resources, limitation of early therapy, or underuse of resources. Finally, this cohort represents ten African countries, despite 26 country leaders agreeing to participate. This result shows the difficulty in fulfilling ethics and regulatory requirements and other barriers associated with doing research in an under-resourced environment.²⁹ It is therefore difficult to determine the generalisability of these results although, to the best of our knowledge, these data provide the largest cohort of critically ill patients with COVID-19 who are from under-resourced environments (appendix p 28).

This is a large, prospective, multicentre study from a previously unreported African setting and, to the best of our knowledge, the only study in this setting that has also included a large number of patients with HIV. The statistical analysis plan was published before data inspection and was adequately powered to adjust for the association between human resources, patient comorbidities, and critical care interventions and mortality. All prespecified sensitivity analyses confirm the main findings.

Hospital and critical care resources are scarce in Africa. Moreover, admission to critical care is restricted and access to critical care interventions are between seven-times and 14-times less than what is needed. Patient comorbidities of HIV/AIDS, diabetes, chronic liver disease and kidney disease, as well as increasing age, are associated with increased mortality from COVID-19 in Africa. Poor prognosis was associated with the degree of organ dysfunction at admission and the need for invasive mechanical ventilation or inotropic support. Although the full SOFA score has been shown to have superior performance to the quick SOFA score,³⁰ most sites analysed in our study could not assess a full SOFA score because of the scarcity of resources. The quick SOFA score is a simple risk stratification or triage tool that is feasible in low resource environments.

Mortality is associated with organ dysfunction and organ support needed at critical admission; yet there are insufficient resources to provide adequate support in this setting. Early warning systems, risk stratification, and early intervention are needed to avoid delays in instituting necessary organ support. Strategies are needed to mitigate risk in patients who are infected with SARS-CoV-2 in Africa with coexisting HIV/AIDS, diabetes, chronic liver disease, and kidney disease.

Contributors

BMB contributed to the overall conception and design of the study, acquisition of data, local and national study leadership, statistical analysis, writing the first draft of the paper, critically revising the work for submission, and the final approval of the version of the study to be submitted. MMi, WLM, DT, AA, EA, GC, HTD, MELi, MG, AGB, IJ, FK, H-LK, ZM, AM, WM, AN, ZN, AO, JLP, JS, YSA, DEAvS, and PDG contributed to the overall conception and design of the study, acquisition of data, local and national study leadership, critical revision of the paper for submission, and were involved in final approval of the study to be submitted. MSC contributed to the overall conception and design of the study, acquisition of data, local and national study leadership, patient recruitment, and data collection. MELh applied for nationwide and local

hospital ethical approvals and recruited collaborators for data collection from the hospital, was involved in local and national study leadership, critically revised the study for submission, and was involved in the final approval of the version of the study to be submitted. MF contributed to the overall conception and design of the study, acquisition of data, local and national study leadership, critical revision of the work for submission, final approval of the version of the study to be submitted, was a national hospital team leader, and was involved in patient recruitment in the study. DF was involved in designing the intensive care unit triage form, acquisition of data, critical revision of the work for submission, and final approval of the version of the study to be submitted. AH was involved in the statistical analyses, critical revision of the work for submission, and the final approval of the version of the study to be submitted. VM and CO were involved in the acquisition of data, local and national study leadership, critical revision of the work for submission, and final approval of the version of the study to be submitted. EHT was involved in the conception and design of the meta-analysis, the acquisition of data for meta-analysis, data analysis and writing, critical revision of the work for submission, and final approval of the version to be submitted. MME contributed to the overall conception and design of the study, local and national study leadership, and the final approval of the version of the study to be submitted. MEI was involved in local and national leadership, critical revision of the work for submission, and final approval of the version of the study to be submitted. KDM was involved in the acquisition of data for meta-analysis, data analysis and writing, critical revision of the work for submission, and final approval of the version to be submitted. FP was involved in designing the intensive care unit triage form, acquisition of data, critical revision of the work for submission, and final approval of the version of the study to be submitted. BMB and AH accessed and verified the data.

Declaration of interests

MME has received honoraria for services related to speakers bureau and advisory boards. These have related to purely educational talks that have been given in an objective fashion for educational purposes and with no vested interest or agenda other than for educational purposes. Companies that MME gave talks to were Pfizer, Merck, Astellas, Sanofi-Aventis, Aspen, and Sun. IJ is the former president of the Critical Care Society of Southern Africa and is a current councillor and board member of the Critical Care Society of Southern Africa. All other authors declare no competing interests.

Data sharing

Data will be disclosed only upon request and approval of the proposed use of the data by the steering committee. Data are available to the journal for evaluation of reported analyses. Data requests from non-ACCOS investigators will not be considered until 2 years after the close of the trial. Data will be de-identified for participant, hospital, and country, and will be available with a signed data access agreement.

Acknowledgments

We would like to acknowledge Rema Ramakrishnan for their assistance with the meta-analysis forest plot and Dilshaad Brey for their assistance with the database search for the meta-analysis. The ACCOS was supported by a grant from the Critical Care Society of Southern Africa.

References

- 1 The Johns Hopkins University School of Medicine. Coronavirus resource center. <https://coronavirus.jhu.edu/map.html> (accessed Sept 30, 2020).
- 2 Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; **395**: 1054–62.
- 3 Biccard BM, Madiba TE, Kluyts HL, et al. Perioperative patient outcomes in the African Surgical Outcomes Study: a 7-day prospective observational cohort study. *Lancet* 2018; **391**: 1589–98.
- 4 Ayebale AET, Kassebaum NJ, Roche AM, Biccard BM. Africa's critical care capacity before COVID-19. *S Afr J Anaesthesiol Analg* 2020; **26**: 162–64.
- 5 Skinner DL, De Vasconcellos K, Wise R, et al. Critical care admission of South African (SA) surgical patients: results of the SA Surgical Outcomes Study. *S Afr Med J* 2017; **107**: 411–19.
- 6 Armstrong RA, Kane AD, Cook TM. Outcomes from intensive care in patients with COVID-19: a systematic review and meta-analysis of observational studies. *Anaesthesia* 2020; **75**: 1340–49.
- 7 Gupta M, Wahl B, Adhikari B, et al. The need for COVID-19 research in low- and middle-income countries. *Glob Health Res Policy* 2020; **5**: 33.
- 8 Taylor EH, Hofmeyr R, Torborg A, et al. Risk factors and interventions associated with mortality or survival in adult COVID-19 patients admitted to critical care: a systematic review and meta-analysis. *S Afr J Anaesthesiol Analg* 2020; **26**: 116–27.
- 9 Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996; **49**: 1373–79.
- 10 Biccard BM, Miller MG, Michell WL, et al. An African, multi-centre evaluation of patient care and clinical outcomes for patients with COVID-19 infection admitted to high-care or intensive care units. *SSRN* 2020; published online Oct 22. https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3707415 (preprint).
- 11 Seymour CW, Liu VX, Iwashyna TJ, et al. Assessment of clinical criteria for sepsis: for the third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016; **315**: 762–74.
- 12 Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the working group on sepsis-related problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996; **22**: 707–10.
- 13 Bates D, Maechler M, Bolker B, Walker SA. Fitting linear mixed-effects models using lme4. *J Stat Softw* 2015; **67**: 1–48.
- 14 RStudio Team. RStudio. Boston, MA: PBC, 2020.
- 15 van Buuren S, Groothuis-Oudshoorn K. mice: multivariate imputation by chained equations in R. *J Stat Softw* 2011; **45**: 1–67.
- 16 Rubin DB. Multiple imputation for nonresponse in surveys. New York: John Wiley & Sons, 1987.
- 17 R Core Team. A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2019.
- 18 Ritchie H, Ortiz-Ospina E, Beltekian D, et al. Mortality risk of COVID-19. 2020. <https://ourworldindata.org/mortality-risk-covid> (accessed Dec 14, 2020).
- 19 Biswas M, Rahaman S, Biswas TK, Haque Z, Ibrahim B. Association of sex, age, and comorbidities with mortality in COVID-19 patients: a systematic review and meta-analysis. *Intervirology* 2021; **64**: 36–47.
- 20 Langer A, Meleis A, Knaut FM, et al. Women and health: the key for sustainable development. *Lancet* 2015; **386**: 1165–210.
- 21 Costenaro P, Minotti C, Barbieri E, Giaquinto C, Donà D. SARS-CoV-2 infection in people living with HIV: a systematic review. *Rev Med Virol* 2021; **31**: 1–12.
- 22 The RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with Covid-19. *N Engl J Med* 2021; **384**: 693–704.
- 23 Nguyen YL, Wallace DJ, Yordanov Y, et al. The volume-outcome relationship in critical care: a systematic review and meta-analysis. *Chest* 2015; **148**: 79–92.
- 24 Hirsch JS, Ng JH, Ross DW, et al. Acute kidney injury in patients hospitalized with COVID-19. *Kidney Int* 2020; **98**: 209–18.
- 25 Combes A, Hajage D, Capellier G, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med* 2018; **378**: 1965–75.
- 26 Tzotzos SJ, Fischer B, Fischer H, Zeitlinger M. Incidence of ARDS and outcomes in hospitalized patients with COVID-19: a global literature survey. *Crit Care* 2020; **24**: 516.
- 27 Barbaro RP, MacLaren G, Boonstra PS, et al. Extracorporeal membrane oxygenation support in COVID-19: an international cohort study of the Extracorporeal Life Support Organization registry. *Lancet* 2020; **396**: 1071–78.
- 28 Romero Starke K, Peterleit-Haack G, Schubert M, et al. The age-related risk of severe outcomes due to COVID-19 infection: a rapid review, meta-analysis, and meta-regression. *Int J Environ Res Public Health* 2020; **17**: E5974.
- 29 Conradie A, Duys R, Forget P, Biccard BM. Barriers to clinical research in Africa: a quantitative and qualitative survey of clinical researchers in 27 African countries. *Br J Anaesth* 2018; **121**: 813–21.
- 30 Liu S, Yao N, Qiu Y, He C. Predictive performance of SOFA and qSOFA for in-hospital mortality in severe novel coronavirus disease. *Am J Emerg Med* 2020; **38**: 2074–80.



Excess COVID-19 mortality among critically ill patients in Africa

See [Articles](#) page 1885

As of March 25, 2021, a global total of 124 215 843 cases of COVID-19, including 2 734 374 deaths, had been reported to WHO.¹ COVID-19 is now the 12th leading cause of death worldwide, the sixth leading cause of death in high-income countries, and the 41st leading cause of death in sub-Saharan Africa.² However, the reasons why there are lower numbers of cases and deaths reported in sub-Saharan Africa are unclear. One possibility is that there are inadequate levels of testing, which could translate to unreported COVID-19 deaths (both in hospital and those that occur outside of hospital).³

A study³ in Zambia detected COVID-19 in 70 (19.2%) of 364 deceased individuals, most of these occurring outside of hospital. In their multicentre, prospective, observational cohort study of 3140 critically ill patients (60.6% male, mean age 55.6 years [SD 16.1]) enrolled from 64 hospitals in ten African countries reported in *The Lancet*, the African COVID-19 Critical Care Outcomes Study (ACCCOS) Investigators⁴ show that, despite having low COVID-19 mortality rates, Africa has the highest global mortality rate in patients with COVID-19 who are critically ill: 48.2% (95% CI 46.4–50.0; 1483 of 3077 patients) against a global average of 31.5% (27.5–35.5). In addition to the previously reported drivers of mortality (eg, the patient's disease severity at presentation and having comorbidities such as HIV/AIDS, diabetes, and chronic liver disease), the ACCCOS Investigators found that having HIV/AIDS (odds ratio 1.91) and delayed access to high-care units and intensive care units (2.14) were drivers of mortality.

Their study is, to the best of our knowledge, the first multicountry report of outcomes of critically ill patients with COVID-19 in Africa. The question is which factors drive this high mortality in a continent with lower cases of COVID-19 and overall lower mortality rates? The authors provide some insights into possible causes, such as a shortage of critical care resources and underuse of those that are available. The underuse of resources is an intriguing finding and contrary to popular belief that resources are scarce. It is shocking to see that 68% of hospitals had access to dialysis but only 10% of the patients received it, as well as to see that proning was not optimised. It is important to think beyond the availability of resources and to also consider issues of functionality. It is common in Africa to have expensive equipment that is non-functional due to poor maintenance or lack of skilled human resources. In 2017, the Tropical Health and Education Trust reported that 40% of the medical equipment in Africa was out of service, 80% of the medical equipment was donated, 70–90% of the donated equipment was never operationalised, and only two African countries had professional biomedical engineers.⁵

The ACCCOS Investigators also report high rates of 24 h physician coverage and a nurse-to-patient ratio of 1:2. Despite this fairly good physician and nurse coverage, mortality was high, possibly because of inadequate skill sets. There are only a small number of pulmonary and critical care training programmes in Africa. Only recently has critical care been added to the anaesthesia curricula in a few African countries. With the exception of Ethiopia, most pulmonary and critical care programmes are in South Africa.⁶ In Tanzania, a survey found that there were 0.04 anaesthetists per 100 000 population and 0.15 anaesthetists of any type per 100 000 population.⁷

The high COVID-19-related mortality in Africa could also be a reflection of the virulence of SARS-CoV-2 as a pathogen. Never before has the world seen a disease that causes the severity of respiratory failure like that caused by SARS-CoV-2. Viruses constantly mutate, leading to variants. Variants of SARS-CoV-2 have recently emerged, including B.1.1.7, a highly transmissible variant that was initially identified in the south of England in September, 2020; P.1, circulating in Brazil since the middle of 2020; and B.1.351, which was



James Oatway/Panos Pictures

first detected in South Africa in late 2020.⁸ The capacity to detect variants in Africa is limited because of inadequate skill and infrastructure for genomic sequencing. The variants have been associated with increased transmissibility and could affect the effectiveness of COVID-19 vaccines. The role of SARS-CoV-2 variants in disease severity is unclear, with only a few reports of increased severity.⁹ Could variants be responsible for the severity seen in this study? This is a question which, in a continent with severe shortage of sequencing, could take a long time to answer.

This study has several strengths, including a large sample size, robust analyses, as well as having a multisite and prospective design. However, the authors also recognise some limitations, including that the study was done in tertiary hospitals. Moreover, 23 (36%) of 64 hospitals were in South Africa and Egypt, which are better resourced countries compared with some other African countries; mortality is probably higher in lower-income African countries. Missing data were overcome by imputation. The authors, however, do not report reasons why one in two patients died without receiving oxygen. Overall, this is a well done study and the team must be congratulated.

We declare no competing interests.

*Bruce J Kirenga, Pauline Byakika-Kibwika
brucekirenga@yahoo.co.uk

Lung Institute (BJK) and Department of Medicine (BJK, PB-K), Makerere University, Kampala, Uganda

- 1 WHO. WHO coronavirus disease (COVID-19) dashboard. <https://covid19.who.int/> (accessed March 25, 2021).
- 2 Institute of Health Metrics and Evaluation. COVID 19 projections—global. 2021. <https://covid19.healthdata.org/global?view=total-deaths&tab=trend> (accessed March 25, 2021).
- 3 Mwananyanda L, MacLeod W, Kwenda G, et al. COVID-19 deaths detected in a systematic post-mortem surveillance study in Africa. *BMJ* 2021; **372**: n334.
- 4 The African COVID-19 Critical Care Outcomes Study (ACCCOS) Investigators. Patient care and clinical outcomes for patients with COVID-19 infection admitted to African high-care or intensive care units (ACCCOS): a multicentre, prospective, observational cohort study. *Lancet* 2021; **397**: 1885–94.
- 5 Worm A, Griffith-Jones L, Namahungu T, et al. The (improved) status of medical equipment in sub-Saharan Africa Healthcare Technology management. May 11, 2017. https://www.who.int/medical_devices/global_forum/3rd_gfmd/statusmedicalequipmentSub-SaharanAfrica.pdf?ua=1 (accessed March 25, 2021).
- 6 Daniel O, Obaseki JEC. Training specialists in developing countries: a new model. *J Pan Afr Thorac Soc* 2020; **2**: 4–5.
- 7 Dubowitz G, Detlefs S, McQueen KAK. Global anesthesia workforce crisis: a preliminary survey revealing shortages contributing to undesirable outcomes and unsafe practices. *World J Surg* 2010; **34**: 438–44.
- 8 Burki T. Understanding variants of SARS-CoV-2. *Lancet* 2021; **397**: 462.
- 9 UK Government. NERVTAG paper on COVID-19 variant of concern B.1.1.7. Jan 22, 2021. <https://www.gov.uk/government/publications/nervtag-paper-on-covid-19-variant-of-concern-b117> (accessed March 25, 2021).